

---

# Pancreatic Neuroendocrine Tumors Causes, Risk Factors, and Prevention

Learn about the risk factors for pancreatic neuroendocrine tumors and what you might be able to do to help lower your risk.

## Risk Factors

A risk factor is anything that affects your chance of getting a disease such as cancer. Learn more about the risk factors for pancreatic neuroendocrine tumors.

- [Pancreatic Neuroendocrine Tumor Risk Factors](#)
- [What Causes Pancreatic Neuroendocrine Tumor?](#)

## Prevention

There is no way to prevent all pancreatic neuroendocrine tumors. But there are things you can do that might lower your risk. Learn more.

- [Can Pancreatic Neuroendocrine Tumor Be Prevented?](#)

---

# Pancreatic Neuroendocrine Tumor Risk Factors

- [Risk factors that can be changed](#)
- [Risk factors that can't be changed](#)
- [Factors with unclear effect on risk](#)

A risk factor is anything that increases your chance of getting a disease such as cancer. Different cancers have different risk factors. Some risk factors, like smoking, can be changed. Others, like a person's age or family history, can't be changed.

But having a risk factor, or even many risk factors, does not mean that you will get the disease. And some people who get the disease may have few or no known risk factors.

Several factors can affect a person's chance of getting a neuroendocrine tumor (NET) of the pancreas.

## **Risk factors that can be changed**

### **Smoking**

[Smoking](#)<sup>1</sup> is a risk factor for pancreatic NETs. Most research shows that heavy smoking increases risk, but some studies show that any history of smoking could put you at risk.

### **Alcohol**

Some studies have shown a link between [heavy alcohol use](#)<sup>2</sup> and pancreatic NETs. This link appears to be mostly related to functioning pancreatic NETs rather than nonfunctioning pancreatic NETs. Heavy alcohol use can also lead to conditions such as chronic pancreatitis, which may increase pancreatic NET risk.

## **Risk factors that can't be changed**

### **Family history**

Pancreatic NETs seem to run in some families. In some of these families, the high risk is due to an inherited syndrome (explained below). In other families, the gene causing the increased risk is not known. If family history is a risk factor, it usually involves a first degree relative (parent, sibling, child), a family history of pancreatic NET, or a family history of any cancer.

### **Inherited genetic syndromes**

Inherited gene changes (mutations) can be passed from parent to child. Sometimes these changes result in syndromes that include increased risks of other cancers (or other health problems).

Pancreatic neuroendocrine tumors and cancers can also be caused by genetic syndromes, such as:

- **Neurofibromatosis, type 1**, which is caused by mutations in the *NF1* gene. This syndrome leads to an increased risk of many tumors, including somatostatinomas.
- **Multiple endocrine neoplasia, type I (MEN1)**, caused by mutations in the *MEN1* gene. This syndrome leads to an increased risk of tumors of the parathyroid gland, the pituitary gland, and the islet cells of the pancreas.
- **Von Hippel-Lindau (VHL) syndrome**, which is caused by mutations in the *VHL* gene. This syndrome leads to an increased risk of many tumors, including pancreatic NETs.

Changes in the genes that cause some of these syndromes can be found by genetic testing. For more information on genetic testing, see [Can Pancreatic Neuroendocrine Tumor Be Found Early?](#)<sup>3</sup>

## Diabetes

Pancreatic NETs are more common in people with diabetes. The reason for this is not known. Most of the risk is found in people with type 2 diabetes. This type of diabetes most often starts in adulthood and is often related to being overweight or obese. It's not clear if people with type 1 (juvenile) diabetes have a higher risk.

## Chronic pancreatitis

Chronic pancreatitis, a long-term inflammation of the pancreas, is linked with an increased risk of pancreatic NETs. If chronic pancreatitis is because of heavy alcohol use, then stopping alcohol may help decrease the risk of pancreatic NETs.

## Factors with unclear effect on risk

### Being overweight or obese

[Being overweight](#)<sup>4</sup> or obese could be a risk factor for pancreatic NET. Studies so far are inconclusive.

## Coffee

Some older studies have suggested that drinking coffee might increase the risk of pancreatic NET, but more recent studies have not confirmed this.

## Hyperlinks

1. [www.cancer.org/cancer/risk-prevention/tobacco.html](http://www.cancer.org/cancer/risk-prevention/tobacco.html)
2. [www.cancer.org/cancer/risk-prevention/diet-physical-activity/alcohol-use-and-cancer.html](http://www.cancer.org/cancer/risk-prevention/diet-physical-activity/alcohol-use-and-cancer.html)
3. [www.cancer.org/cancer/types/pancreatic-neuroendocrine-tumor/detection-diagnosis-staging/detection.html](http://www.cancer.org/cancer/types/pancreatic-neuroendocrine-tumor/detection-diagnosis-staging/detection.html)
4. [www.cancer.org/cancer/risk-prevention/diet-physical-activity/body-weight-and-cancer-risk.html](http://www.cancer.org/cancer/risk-prevention/diet-physical-activity/body-weight-and-cancer-risk.html)

## References

Ben Q, Zhong J, Fei J, et al. Risk Factors for Sporadic Pancreatic Neuroendocrine Tumors: A Case-Control Study. *Scientific Reports*. 2016;6:36073. doi:10.1038/srep36073.

Cancer.Net. Von Hippel-Lindau Syndrome. 12/2017. Accessed at <https://www.cancer.net/cancer-types/von-hippel-lindau-syndrome> on October 8, 2018.

Haugvik SP, Hedenström P, Korsæth E, et al. Diabetes, smoking, alcohol use, and family history of cancer as risk factors for pancreatic neuroendocrine tumors: a systematic review and meta-analysis. *Neuroendocrinology*. 2015;101(2):133-42. doi: 10.1159/000375164. Epub 2015 Jan 22.

Leoncini E, Carioli G, La Vecchia C, Boccia S, Rindi G. Risk factors for neuroendocrine neoplasms: a systematic review and meta-analysis. *Ann Oncol*. 2016 Jan;27(1):68-81. doi: 10.1093/annonc/mdv505. Epub 2015 Oct 20.

National Cancer Institute. Physician Data Query (PDQ). Pancreatic Neuroendocrine Tumors (Islet Cell Tumors) Treatment – Patient Version. 2018. Accessed at <https://www.cancer.gov/types/pancreatic/patient/pnet-treatment-pdq> on October 7, 2018.

Valente R, Hayes AJ, Haugvik SP, et al. Risk and protective factors for the occurrence

of sporadic pancreatic endocrine neoplasms. *Endocr Relat Cancer*. 2017 Aug;24(8):405-414. doi: 10.1530/ERC-17-0040. Epub 2017 May 31.

Vinik A, Perry RR, Hughes MS, et al. Multiple Endocrine Neoplasia Type 1. [Updated 2017 Oct 7]. In: De Groot LJ, Chrousos G, Dungan K, et al., editors. Endotext [Internet]. South Dartmouth (MA): MDText.com, Inc.; 2000-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK481897/>. Accessed October 9, 2018.

Last Revised: October 30, 2018

## What Causes Pancreatic Neuroendocrine Tumor?

- [Inherited gene mutations](#)
- [Acquired gene mutations](#)

Scientists don't know exactly what causes most pancreatic neuroendocrine tumors (NETs), but they have found several [risk factors](#) that can make a person more likely to get this disease. Some of these risk factors affect the DNA of cells in the neuroendocrine system in the pancreas, which can result in abnormal cell growth and may cause cancers to form.

DNA is the chemical in our cells that carries our genes, which control how our cells function. We look like our parents because they are the source of our DNA. But DNA affects more than just how we look.

Some [genes](#)<sup>1</sup> control when our cells grow, divide into new cells, and die:

- Genes that help cells grow, divide, and stay alive are called *oncogenes*.
- Genes that help keep cell division under control, repair mistakes in DNA, or cause cells to die at the right time are called *tumor suppressor genes*.

Cancers can be caused by DNA changes (mutations) that turn on oncogenes or turn off tumor suppressor genes.

## Inherited gene mutations

Although 90% of PNETs are sporadic (random), some people [inherit gene changes](#)<sup>2</sup> from their parents that raise their risk of pancreatic NET. Sometimes these gene changes are part of syndromes that include increased risks of other health problems as well.

Syndromes related to changes in three tumor suppressor genes are responsible for many inherited cases of pancreatic NETs:

- **Multiple Endocrine Neoplasia Type 1 (MEN1) syndrome:** Most inherited cases of PNETs are due to changes in the MEN1 gene. This syndrome can cause cancer in the pancreas, parathyroid glands, and [pituitary glands](#)<sup>3</sup>. These tumors usually happen at younger ages and tend to be non-functioning. Screening people with the MEN1 gene or their family members can sometimes help find pancreatic NET before symptoms appear.
- **Von Hippel-Lindau (VHL) syndrome:** Changes in the *VHL* gene cause a small number of pancreatic NETs, usually developing at earlier ages (sometimes as early as the 20s). These tumors tend to be non-functioning and slow growing.
- **Neurofibromatosis type 1 (NF1) syndrome:** A small number of pancreatic NETs (usually somatostatinomas) are caused by changes in the *NF1* gene. Other cancers are also associated with this syndrome, including brain tumors or benign tumors that form in nerves under the skin (neurofibromas),

The treatment for a pancreatic NET that's caused by a genetic syndrome might be slightly different compared to treatment for a pancreatic NET in someone without a gene mutation.

## Acquired gene mutations

Most gene mutations related to neuroendocrine tumors of the pancreas are caused by random changes. These random mutations are called *acquired* if they occur after a person is born, rather than having been inherited. These acquired gene mutations sometimes result from exposure to cancer-causing chemicals (like those found in [tobacco smoke](#)<sup>4</sup>). But often what causes these changes is not known.

## Hyperlinks

1. [www.cancer.org/cancer/risk-prevention/genetics.html](http://www.cancer.org/cancer/risk-prevention/genetics.html)
2. [www.cancer.org/cancer/risk-prevention/genetics.html](http://www.cancer.org/cancer/risk-prevention/genetics.html)
3. [www.cancer.org/cancer/types/pituitary-tumors.html](http://www.cancer.org/cancer/types/pituitary-tumors.html)
4. [www.cancer.org/cancer/risk-prevention/tobacco/carcinogens-found-in-tobacco-products.html](http://www.cancer.org/cancer/risk-prevention/tobacco/carcinogens-found-in-tobacco-products.html)

## References

Benafif S and Eeles R. Diagnosis and Management of Hereditary Carcinoids. *Recent Results Cancer Res.* 2016; 205:149-68. doi: 10.1007/978-3-319-29998-3\_9.

Chai SM, Brown IS, Kumarasinghe MP. Gastroenteropancreatic neuroendocrine neoplasms: selected pathology review and molecular updates. *Histopathology.* 2018 Jan;72(1):153-167. doi: 10.1111/his.13367.

Chen M, Van Ness M, Guo Y, Gregg J. Molecular pathology of pancreatic neuroendocrine tumors. *Journal of Gastrointestinal Oncology.* 2012;3(3):182-188. doi:10.3978/j.issn.2078-6891.2012.018.

Friedman JM. Neurofibromatosis 1. 1998 Oct 2 [Updated 2018 May 17]. In: Adam MP, Ardinger HH, Pagon RA, et al., editors. GeneReviews® [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2018. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK1109/>

Lee J, Allendorf J, Chabot J. Surgical resection of sporadic pancreatic neuroendocrine tumors. UpToDate website. <https://www.uptodate.com/contents/surgical-resection-of-sporadic-pancreatic-neuroendocrine-tumors>. Updated January 09, 2018. Accessed October 3, 2018.

van Leeuwen RS, Ahmad S, Links TP, et al. Von Hippel-Lindau Syndrome. 2000 May 17 [Updated 2018 Sep 6]. In: Adam MP, Ardinger HH, Pagon RA, et al., editors. GeneReviews® [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2018. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK1463/>

Vinik A, Perry RR, Hughes MS, et al. Multiple Endocrine Neoplasia Type 1. [Updated 2017 Oct 7]. In: De Groot LJ, Chrousos G, Dungan K, et al., editors. Endotext [Internet]. South Dartmouth (MA): MDText.com, Inc.; 2000-. Available from:

<https://www.ncbi.nlm.nih.gov/books/NBK481897/>. Accessed 10.02.18.

Yao JC, Evans DB. Chapter 85: Pancreatic neuroendocrine tumors. In: DeVita VT, Lawrence TS, Rosenberg SA, eds. *DeVita, Hellman, and Rosenberg's Cancer: Principles and Practice of Oncology*. 10th ed. Philadelphia, Pa: Lippincott Williams & Wilkins; 2015.

Last Revised: October 30, 2018

## Can Pancreatic Neuroendocrine Tumor Be Prevented?

- [Don't smoke](#)
- [Limit alcohol use](#)

There is no sure way to prevent pancreatic neuroendocrine tumors (NETs). Some [risk factors](#) such as family history can't be controlled. But there are things you can do that might lower your risk.

### Don't smoke

Smoking is an avoidable risk factor for pancreatic NET. [Quitting smoking](#)<sup>1</sup> may help lower risk. If you smoke and want help quitting, please talk to your health care provider or call us at 1-800-227-2345.

### Limit alcohol use

Heavy alcohol use has been tied to pancreatic NETs in some studies but not in others. This link is still not certain, but heavy alcohol use can lead to conditions such as chronic pancreatitis, which has been associated with an increased risk of pancreatic NET.

### [Hyperlinks](#)



1. [www.cancer.org/cancer/risk-prevention/tobacco.html](http://www.cancer.org/cancer/risk-prevention/tobacco.html)

## References

Ben Q, Zhong J, Fei J, et al. Risk Factors for Sporadic Pancreatic Neuroendocrine Tumors: A Case-Control Study. *Scientific Reports*. 2016;6:36073. doi:10.1038/srep36073.

Haugvik SP, Hedenström P, Korsæth E, et al. Diabetes, smoking, alcohol use, and family history of cancer as risk factors for pancreatic neuroendocrine tumors: a systematic review and meta-analysis. *Neuroendocrinology*. 2015;101(2):133-42. doi: 10.1159/000375164. Epub 2015 Jan 22.

Leoncini E, Carioli G, La Vecchia C, Boccia S, Rindi G. Risk factors for neuroendocrine neoplasms: a systematic review and meta-analysis. *Ann Oncol*. 2016 Jan;27(1):68-81. doi: 10.1093/annonc/mdv505. Epub 2015 Oct 20.

Last Revised: October 30, 2018

## Written by

The American Cancer Society medical and editorial content team  
(<https://www.cancer.org/cancer/acs-medical-content-and-news-staff.html>)

Our team is made up of doctors and oncology certified nurses with deep knowledge of cancer care as well as journalists, editors, and translators with extensive experience in medical writing.

American Cancer Society medical information is copyrighted material. For reprint requests, please see our Content Usage Policy ([www.cancer.org/about-us/policies/content-usage.html](http://www.cancer.org/about-us/policies/content-usage.html)).

**cancer.org | 1.800.227.2345**