

Incidence and Mortality Rates for Selected Cancers by Race and Ethnicity, US



| Incidence, 2016-2020 | All races & ethnicities | White | Black | American Indian/ Alaskan Native† | Asian American/ Pacific Islander | Hispanic/ Latino |
|--------------------------------|-------------------------|-------|-------|-------------------------------------|-------------------------------------|---------------------|
| All sites | 453.2 | 474.3 | 459.7 | 478.8 | 301.3 | 358.1 |
| Male | 492.5 | 511.2 | 533.9 | 504.1 | 299.0 | 377.2 |
| Female | 426.6 | 449.3 | 409.9 | 465.5 | 307.3 | 351.3 |
| Breast (female) | 129.0 | 134.9 | 129.6 | 115.5 | 104.6 | 100.7 |
| Colon & rectum* | 35.3 | 35.2 | 40.8 | 50 | 28.1 | 32.2 |
| Male | 40.7 | 40.4 | 48.8 | 57.8 | 33.4 | 38.2 |
| Female | 30.6 | 30.5 | 35.0 | 43.7 | 23.7 | 27.2 |
| Kidney & renal pelvis | 17.6 | 17.8 | 19.3 | 33.0 | 8.2 | 17.9 |
| Male | 23.9 | 24.3 | 26.4 | 43.9 | 11.6 | 23.5 |
| Female | 12.1 | 12.1 | 13.7 | 23.9 | 5.5 | 13.3 |
| Liver & intrahepatic bile duct | 8.8 | 7.5 | 10.5 | 19.1 | 11.9 | 13.9 |
| Male | 13.2 | 11.2 | 17.0 | 27.3 | 18.4 | 20.4 |
| Female | 4.9 | 4.2 | 5.5 | 12.3 | 6.7 | 8.4 |
| Lung & bronchus | 55.0 | 59.5 | 56.7 | 62.2 | 33.6 | 28.3 |
| Male | 62.2 | 65.7 | 72.4 | 67.2 | 40.8 | 34.3 |
| Female | 49.4 | 54.8 | 45.8 | 58.6 | 28.1 | 24.0 |
| Prostate | 115.0 | 110.7 | 186.1 | 91.9 | 60.9 | 90.9 |
| Stomach | 6.3 | 5.1 | 9.7 | 10.1 | 9.0 | 9.3 |
| Male | 8.4 | 7.1 | 13.0 | 13.1 | 11.8 | 11.4 |
| Female | 4.6 | 3.4 | 7.4 | 7.8 | 6.9 | 7.7 |
| Uterine cervix | 7.7 | 7.2 | 8.6 | 11.4 | 6.0 | 9.7 |

Mortality, 2016-2020

| | | | | | | |
|--------------------------------|-------|-------|-------|-------|-------|-------|
| All sites | 149.8 | 155.0 | 175.8 | 183.8 | 95.4 | 108.6 |
| Male | 178.0 | 183.3 | 217.4 | 221.6 | 111.6 | 130.2 |
| Female | 129.1 | 133.6 | 150.2 | 157.9 | 83.7 | 93.5 |
| Breast (female) | 19.7 | 19.7 | 27.8 | 21.1 | 11.8 | 13.7 |
| Colon & rectum | 13.2 | 13.1 | 17.7 | 19.0 | 9.2 | 10.7 |
| Male | 15.7 | 15.5 | 22.4 | 23.1 | 11.0 | 13.6 |
| Female | 11.0 | 11.1 | 14.4 | 16.0 | 7.8 | 8.5 |
| Kidney & renal pelvis | 3.5 | 3.6 | 3.4 | 6.7 | 1.6 | 3.3 |
| Male | 5.1 | 5.3 | 5.2 | 9.9 | 2.4 | 4.8 |
| Female | 2.2 | 2.3 | 2.2 | 4.2 | 1.0 | 2.1 |
| Liver & intrahepatic bile duct | 6.6 | 5.9 | 8.3 | 13.6 | 8.5 | 9.3 |
| Male | 9.6 | 8.5 | 13.0 | 19.9 | 12.6 | 13.1 |
| Female | 4.1 | 3.7 | 4.8 | 8.8 | 5.2 | 6.0 |
| Lung & bronchus | 35.0 | 38.2 | 37.5 | 43.4 | 20.0 | 15.5 |
| Male | 42.3 | 44.9 | 51.3 | 52.3 | 25.9 | 21.0 |
| Female | 29.4 | 32.9 | 28.0 | 37.0 | 15.6 | 11.4 |
| Prostate | 18.9 | 17.9 | 37.9 | 22.5 | 8.7 | 15.4 |
| Stomach | 2.9 | 2.1 | 5.0 | 5.6 | 4.7 | 4.8 |
| Male | 3.8 | 2.9 | 7.2 | 7.7 | 6.0 | 5.9 |
| Female | 2.1 | 1.5 | 3.5 | 4.1 | 3.7 | 3.9 |
| Uterine cervix | 2.2 | 2.0 | 3.3 | 3.3 | 1.7 | 2.5 |

Rates are per 100,000 population and age adjusted to the 2000 US standard population; incidence is adjusted for delays in reporting. More information on state exclusion can be found in the Sources of Statistics, page 79. All race groups are exclusive of Hispanic origin (see Sources of Statistics, page 79). *Colorectal cancer incidence rates exclude appendix. †To reduce racial misclassification for American Indians and Alaska Native (AIAN) individuals, incidence rates are limited to Purchased/Referred Care Delivery Area counties. Mortality rates for AIAN individuals are for the entire US and are adjusted for racial misclassification on death certificates using factors from the National Center for Health Statistics (see Sources of Statistics, page 79).

Source: Incidence – North American Association of Central Cancer Registries (NAACCR), 2023. Mortality – National Center for Health Statistics, Centers for Disease Control and Prevention, 2023.